arXiv:1207.2802v1 [physics.soc-ph] 11 Jul 2012

Coupled dynamics of mobility and pattern formation in optional public goods games

Li-Xin Zhong[a], Wen-Juan Xu[a], Yong-Dong Shi[a,b], Tian Qiu[c]

[a]School of Finance, Zhejiang University of Finance and economics, Hangzhou, 310018, China
[b]School of Finance and Research Center of Applied Finance, Dongbei University of Finance and Economics, Dalian, 116025, China
[c]School of Information Engineering, Nanchang Hangkong University, Nanchang, 330063, China

---

Abstract

In a static environment, optional participation and cooperator agglomeration are found to be beneficial for the occurrence and maintenance of cooperation. In the optional public goods game, the rock-scissors-paper cycles of different strategies yield the oscillatory persistence of cooperation but not different levels of stable cooperation. In this paper, by incorporating population density and individual mobility into the spatial optional public goods game, we study the coevolutionary dynamics of strategy updating and benefit-seeking movement. With low population density and slow movement, an optimal level of cooperation is easy to be reached. Both the increase of population density and the speed-up of free-floating of competing agents will lead to the decrease of cooperation. A log-log relation between the levels of cooperation and the free-floating probability is found. Theoretical analysis indicates that the decrease of cooperator frequency in the present model should result from the increase of the interactions between different agents, which may originate from the enlargement of the average size of the interaction groups or the rise of the speed of random-movement.

Keywords: mobility, cooperation, population density, public goods games

---

Email address: `zlxxwj@163.com` (Li-Xin Zhong[a])

1. Introduction

Social dilemmas describe a conflict situation existing between a rational individual maximizing its own benefit and a social group pursuing collective wellbeing[1, 2]. For example, as a sheepherder enjoys herding in the common grass the joint effort to attain a sustainable environment suffers a risk. The gas emission of a factory ensuring its prosperity makes the greenhouse problem become more serious, which in turn do harm to the further development of the factory. As everyone faces the temptation to exploit the public goods and make no contribution to it, the immense benefit that can only be got by all the people's cooperation becomes unattainable. This poses a challenging problem about how the individual selfishness can lead to the occurrence and maintenance of cooperation commonly found in reality? To answer this question, various mechanisms have been introduced by scientists since Darwin[3, 4].

Most studies on the evolution of cooperation among selfish individuals are based on game models, the Prisoner's Dilemma game(PDG)[5, 6, 7, 8], the snowdrift game (SG)[9, 10, 11, 12, 13], and the public goods game (PGG)[14, 15, 16], to name just a few. The iterated PDG models transactions between two agents, in which one's contribution favors the other but not itself. Although the total income would be the highest if they both cooperate, an agent tends to defect to maximize its own profit. Therefore, the Nash equilibrium is to defect at all times. The PGG is an extension of the PD to an arbitrary number of agents. In the original PGG, a group of individuals have the opportunity to make an investment into a common pool or not. An even distribution of returns irrespective of one's contribution makes the defectors have an advantage over the cooperators and defection becomes a dominating strategy in such an evolutionary system. To refrain from getting stuck in the deadlocks of mutual defection, a third strategy, termed a loner's strategy[17, 18, 19, 20], has been introduced into the original public goods game.

In the public goods game with loners, also termed optional public goods game (OPGG), the agents have the option to participate in the public goods game or not. Those who join the public goods game get a cooperator's payoff P_C or a defector's payoff P_D, and those who do not join the public goods game get a loner's payoff P_L. Because a loner only gets a small but fixed payoff, it can win over a group of defectors but will be defeated by a group of cooperators. Therefore, an endless rock-scissors-paper cycle occurs: Loners will invade a population of defectors with fewer cooperators. Cooperators will thrive in a population of loners. A population of cooperators will be intruded into by defectors.

In the optional public goods game, although the existence of loners help the cooperators keep from being doomed, the oscillation of the levels of the three strategies indicates that a higher and stable level of cooperation is difficult to be reached in such a system. But in the real world, cooperation often becomes a dominant strategy in animal and human activities and an environmental change will lead to the occurrence of different levels of cooperation. The environmental conditions include the structured space[21, 22, 23, 24, 25, 26], the population density[27, 28, 29, 30], and the mobility of the individuals[31, 32, 33, 34]. To find out the mechanisms determining the occurrence of different levels of cooperation in the real world, it is a real need to generalize the model presented by Hauert et al. and incorporate the environmental conditions into the original OPGG [17].

In natural and human society, the structured space is not full-occupied and random and purposive movements often occur[35, 36, 37**?**]. Such as the migration of birds, the floating of a boat and the movement of a train. Z.H.Liu has studied the influence of population density and individual mobility on epidemic spreading[39]. C.P.Roca et al and J.Y.Wakano et al have studied the roles of mobility in the improvement of cooperation in the PGG and in the ecological PGG respectively[40, 30]. Motivated by the work done in partially occupied rigid networks, in this paper, we incorporate population density and individual mobility into the original OPGG introduced in ref.[17] and play the game in a square lattice with the Moore neighborhood. The main findings of the study are as follows:

(1) With a predefined free-floating probability ξ_0, the cooperator frequency f_C is determined by population density ρ. There exists a transition point ρ_{tr}, below which f_C increases with the rise of ρ while above which f_C decreases with the rise of ρ. With a predefined ρ, f_C is determined by ξ_0. Increasing ξ_0 leads to a monotonic decrease of f_C.

(2) Considering the size distribution $P(S)$ of individual components, we find that the power-law relation between f_C and ξ_0 is related to the occurrence of the largest component. Before the largest component $S_{max} \sim N$ occurs, as ρ increases, $P(S)$ changes from an exponent to a power-law and the slop of f_C as a function of ξ_0 decreases with the rise of ρ. After the largest component $S_{max} \sim N$ occurs, as ρ increases, $P(S)$ changes little and the slop of f_C as a function of ξ_0 has no change with the rise of ρ.

(3) Taking an eye of the decrease of cooperation in the present model, the effect of increasing ξ_0 with a fixed ρ is similar to the effect of increasing ρ with a fixed ξ_0. The theoretical analysis shows that the change of the frequencies of different strategies in the present model should be determined by the possible collisions of the agents with different strategies. Both the increase of ρ and the increase of ξ_0 would result in more such collisions. The more collisions of the agents, the lower cooperative levels of the system.

The paper is proceeded as follows. In Section 2, we introduce the optional public goods game with purposive and random movements in a spatial setting. In Section 3, simulation results about the evolution of competing strategies and the state of local agglomeration are presented and the relationship between them is discussed. In Section 4, the extinction thresholds are analyzed with mean field theory, and the possible relations between the levels of cooperation and the individual collisions are described theoretically. Section 5 summarizes the paper and gives an outlook of future studies.

2. The model

A population of N agents is distributed over a square lattice with side length L and the Moore neighborhood (i.e., degree $k = 8$), each agent on each site. For $N \leq L^2$, the population density is defined as $\rho = \frac{N}{L^2}$. For $\rho < 1$, an agent can move around and occupy the firstly found empty site. In the present model, there exist two co-evolutionary processes: the change of personal strategies and the movement of the agents. Once the initial position and the adopted strategy of each agent are set, the system can evolve as follows.

In the evolution of personal strategies. Initially, there exist three kinds of agents: cooperators (C), defectors (D) and loners (L). At each time step, firstly, each agent interacts with its neighbors and get a payoff P_C (for cooperators), P_D (for defectors) or P_L (for loners). Owing to the neighboring restriction, the interaction group in the present model should be $n \leq 9$. Assuming $n = n_C + n_D + n_L$, in which n_C, n_D, n_L represent the number of cooperators, defectors and loners in the interaction group respectively, the payoffs for the agents with different strategies are

$$P_C = \frac{rn_C}{n_C + n_D} - 1, \tag{1}$$

$$P_D = \frac{rn_C}{n_C + n_D}, \tag{2}$$

$$P_L = \sigma, \tag{3}$$

in which r (> 1) is the multiplication rate and σ generally satisfies $0 < \sigma < r - 1$, which keeps a loner's payoff to be lower than the payoff of the agents in a group of cooperators and higher than the payoff of the agents in a group of defectors. If $n_C + n_D = 1$, the cooperator or defector will get a loner's payoff. After all the agents have attained their payoffs, they will make decisions on whether they should update their strategies or not. In the updating process, an agent i compares its payoff with a randomly chosen neighbor j's and adopt j's strategy with probability

$$\Gamma_{i \leftarrow j} = \frac{1}{1 + e^{(P_i - P_j + \tau)/\kappa}}, \tag{4}$$

in which τ and κ represent the cost of strategy change and the environmental noise respectively.

The above updating mechanism shows that, in an interaction group, the evolution of the mixed strategy profile is determined by the scores of different strategies. Even if the group size is constant, the number of competitive agents, i.e. cooperators and defectors, will vary with time. Such that the oscillations of the population composition will occur in a fully-occupied network setting.

In relation to the mobility of individuals. For a loner, because it only relies on a small but fixed payoff and gets nothing from a benefit-seeking competition, it is not so attractive to stay at its current site and at each time step it will move randomly with probability ξ_L. For cooperators and defectors, they make their decisions on whether they should leave their current sits or not based on their satisfaction to the present situation and the possible free-floating, which are described by the purposive-moving probability ξ_g and the free-floating probability ξ_0 respectively. The value of ξ_0 is predefined and the value of ξ_g is determined by the wealth of an agent, which is an accumulated payoff in the latest T time steps. For an agent i, its wealth is

$$w_i(t) = \sum_{t'=t-T+1}^{t} p_i(t'). \tag{5}$$

For a cooperator or a defector, if its wealth w is less than 0, with probability

$$\xi_{CD} = \xi_g + \xi_0, \tag{6}$$



Figure 1: Frequencies of cooperators (circles), defectors (squares), and loners (triangles) as a function of multiplication rate r in a square lattice with size $M = L \times L = 200 \times 200$ and Moore neighborhood. Other parameters are $\sigma = 1$, $\tau = \kappa = 0.1$, $\xi_L = 0.1$, $\xi_0 = 0.01$, $\varepsilon_L = 0.001$ and (a)$\rho = 0.3$, (b)$\rho = 0.5$, (c)$\rho = 0.8$. The results are obtained by averaging over 10 runs and 1000 times after 10000 relaxation times in each run.

in which $\xi_g = 1 - e^w$, it leaves its current site and finds an empty site to stay on.

Therefore, in the present model, there are two mechanisms which determine the evolution of cooperation. The payoff determines the replicator dynamics while the wealth determines the mobility dynamics. The great concern of us is how the evolutionary dynamics in a three-strategy game are affected by the population density and the free-floating probability. In the present model, to refrain from the random disappearance of loners, at every time step, the competitive agents will occasionally become loners with a small random-flapping probability ε_L.

3. Results and discussions

In this section, we will focus on the roles of population density and free-floating probability of cooperators and defectors in the change of cooperation. Following the work done in ref.[17], in Monte Carlo simulations we choose the loner's payoff $\sigma = 1$, the cost of strategy change $\tau = 0.1$ and the environmental noise $\kappa = 0.1$. Throughout the paper, the following parameters are also predefined: the size of the square lattice $M = L \times L = 200 \times 200$, the random-moving probability of loners $\xi_L = 0.1$ and the random-flapping $C \to L$ (or $D \to L$) probability $\varepsilon_L = 0.001$.

Figure 1 shows the frequencies of cooperators, defectors and loners as a function of r in a square lattice with different population densities $\rho = 0.3$,



Figure 2: Frequencies of cooperators (circles), defectors (squares), and loners (triangles) as a function of multiplication rate r in a square lattice with $\rho = 0.3$ and (a)$\xi_0 = 0.01$, (b)$\xi_0 = 0.05$, (c)$\xi_0 = 0.1$. All the other parameters are the same as those in fig.1.

0.5 and 0.8. As what has been found in a fully-occupied regular network[25], there exist two extinction thresholds r_D and r_L. For $r < r_D$, loners perform better than cooperators and defectors so that all the agents become loners in the final steady state. For $r_D < r < r_L$, the three strategies coexist. The rise of ρ is found to be beneficial for defectors and loners but not for cooperators. For $r > r_L$, the survival of cooperators is favored and the loners go into extinction.

In all the three cases with different ρ, the values of the extinction threshold r_D are the same, $r_D = \sigma + 1 = 2$, which is also the same as that in a fully-occupied regular network[41]. Such a result comes from the fact that, with a small multiplication rate $r < 2$, even if all the competing agents are cooperators, the payoff of them, $P_C = r - 1$, is less than a loner's payoff $\sigma = 1$. Therefore, the rock-scissors-paper cycle does not occur and the system is stuck in the state where all the agents are loners. However, as we consider the extinction threshold r_L, we find it increases with the rise of ρ. For $\rho = 0.3$, $r_L \sim 4$. For $\rho = 0.5$, $r_L \sim 5.5$. For $\rho = 0.8$, $r_L \sim 6.5$.

The rise of population density can effectively leads to the rise of the probabilities of immediate interactions between two agents. Such an increase is more possible to occur in the system where the agents can move freely. In fig.2, we give the frequencies of different strategies as a function of r with predefined population density $\rho = 0.3$ and different ξ_0. Comparing the results in fig.2 with the results in fig.1, we find that, as to the change of the frequencies of different strategies, the effect of increasing ξ_0 for fixed ρ is similar to the effect of increasing ρ for fixed $\xi_0 (\neq 0)$. As ξ_0 increases, the



Figure 3: Frequencies of (a) cooperators, (b) defectors, and (c) loners as a function of population density ρ for $r = 2.5$ and different $\xi_0 = 0.01$(circles), 0.05(squares), 0.1(triangles). All the other parameters are the same as those in fig.1.

cooperator frequency f_C decreases while the defector frequency f_D and the loner frequency f_L increase within a large range of r. As ξ_0 increases from 0.01 through 0.05 to 0.1, the extinction threshold r_L increases from 4 through 5 to 5.5 accordingly.

The results in fig.1 and fig.2 indicate that, the increase of the meeting probability, whether it comes from the rise of population density or the rise of free-floating probability, will lead to the decrease of cooperation.

It is instructive to ask: can we draw a conclusion that the fewer the chances to meet, the higher the levels of cooperation in the present model? To examine whether an optimal level of population density exists, in fig. 3



Figure 4: Time-dependent frequencies of cooperators with r=2.5, $\xi_L = 0.1$, $\varepsilon_L = 0.001$ and (a) $\xi_0 = 0.01$, $\rho = 0.07$(black), 0.10(red), 0.25(blue); (b)$\xi_0 = 0.01$, $\rho = 0.30$(black), 0.35(red), 0.60(blue); (c) $\rho = 0.3$, $\xi_0 = 0.01$(black), 0.05(red), 0.10(blue).

we give the frequencies of the three strategies as a function of ρ for different ξ_0. With a small and fixed $\xi_0 = 0.01$, there exists a transition point ρ_{tr}. For $\rho < \rho_{tr}$, the defectors do not exist in the population. The cooperators increase while the loners decrease with the rise of ρ. For $\rho > \rho_{tr}$, the cooperators decrease while the loners increase with the rise of ρ. Within this range, the defectors firstly increase and then decrease with the rise of ρ. Increasing the free-floating probability leads to the decrease of the levels of cooperation but not the disappearance of the transition point. As ξ_0 ranges from 0.01 to 0.1, the transition point ρ_{tr} changes from 0.28 to 0.1 and the cooperator frequency at the the transition point decreases from 1 to 0.86 accordingly. The existence of the transition point indicates that, there exists an optimal level of population density, with which the system will reach the highest level of cooperation.

To have a close eye of the evolution of cooperation below and above the transition point ρ_{tr}, in fig. 4(a) and (b) we plot the time-dependent frequencies of cooperators for different ρ. For comparison, we also plot the time-dependent cooperator frequencies for different ξ_0 in fig. 4(c). For $\rho < \rho_{tr}$, as the time passes f_C becomes stable. The change of ρ only leads to the change of the average value of cooperator frequencies but not the fluctuations of it. For $\rho > \rho_{tr}$, the change of ρ not only results in the change of f_C but also the stableness of cooperation. Increasing ρ leads to large fluctuations of the levels of cooperation. Figure 4(c) shows increasing ξ_0 also leads to large fluctuations of the levels of cooperation. Such results indicate that, whether the decrease of cooperator frequency results from the increase of ρ



Figure 5: The size distribution of individual components in the evolved system with r=2.5, $\xi_L = 0.1$, $\varepsilon_L = 0.001$ and different population density $\rho = 0.3$(circles), 0.4(squares), 0.5(triangles).



Figure 6: The size of the largest component as a function of population density ρ in the evolved system with r=2.5, $\xi_L = 0.1$ and $\varepsilon_L = 0.001$.

or the increase of ξ_0, the large fluctuations of the strategies are detrimental to cooperation.

To find the relationship between the frequencies of strategies and the population patterns, in fig.5 we plot the size distribution of individual components for different ρ. As ρ increases from 0.3 through 0.4 to 0.5, the size distribution changes from an exponential distribution through a power-law distribution to a single quite large component accompanied by a power-law distribution.

In fig.6 we give the size of the largest component as a function of ρ. As ρ ranges from 0.1 to 0.3, the size of the largest component has little change with the rise of ρ. As ρ ranges from 0.3 to 0.5, the size of the largest component



Figure 7: Log-log plot of the cooperator frequency f_C vs ξ_0 with $\rho = 0.3$, 0.4, 0.5, 0.6, 0.7. All the other parameters are r=2.5, $\xi_L = 0.1$ and $\varepsilon_L = 0.001$. The fitted lines satisfy the equation $f_C = a\xi_0^{-b}$.

has a sharp increase with the rise of ρ. For $\rho > 0.5$, nearly all the agents are in the same component and the largest component has little change with the rise of ρ.

Comparing the results in fig.3 with the results in fig.5 and fig.6 we find that, for low population density, the agents are scattered and the levels of cooperation are somewhat high. For high population density, nearly all the agents are in the same component and the the levels of cooperation become lower. Therefore, the results in fig.3 can be understood as follows: For quite low population density, the leave of the unsatisfactory agents makes it become impossible for the defectors to exploit the cooperators and the group of defectors will finally be doomed by the random-moving loners. And for the cooperators who leave the competing group, because of the low population density, it is not easy for them to find another cooperator to collaborate with and they will finally become solitary agents whose payoffs are equal to the payoff of a loner. The occasional $C \to L$ (or $D \to L$) random-flapping can make the loners refrain from extinction. Therefore, under such circumstances, only cooperators and loners are left in the final steady state. But for high population density or a high free-floating probability, it is easy for defectors to intrude into cooperator clusters and the average level of cooperation will accordingly decrease with the rise of ρ or the rise of ξ_0.

To have a deep understanding of the roles of free-floating in the increase or decrease of cooperation, in fig.7 we plot f_C vs ξ_0 for different ρ. From fig.7 we find that, for low population density, the frequency of cooperators decreases sharply with the rise of ξ_0. The rise of ρ makes such a changing tendency become ease. As we fit curves to data points in fig.7, it is found that f_C and ξ_0 satisfy the equation $f_C = a\xi_0^{-b}$, in which a=0.28769, 0.37352, 0.39333, 0.37861, 0.34281 and b=0.23855, 0.12845, 0.087778, 0.07006, 0.069364 respectively.

Comparing the results in fig.7 with the results in fig.6, we find that the slope of the fitted line in fig.7 is closely related to the occurrence of the largest component. As the size of the largest component S_{max} is quite small, the slope of the fitted line is steep, which indicates that the change of ξ_0 can quite effectively affect f_C as the agents are scattered. As S_{max} has a sharp increase, the slope of the fitted line quickly becomes gentle. As $S_{max} \sim N$, the change of ρ has little effect on the change of S_{max}, and accordingly the slope of the fitted line no longer changes with the rise of ρ. Such results indicate that, in the present model, the value of b in equation $f_C = a\xi_0^{-b}$ contains the information of evolutionary patterns. We may effectively figure out the size distribution of the components from the slope of the fitted line of $f_C(\xi_0)$.

The above simulation results indicate that, as to the decrease of cooperation, the role of increasing ξ_0 is similar to the effect of increasing ρ. The change of cooperation in the present model should come from the change of the meeting probability of the agents. Both increasing ξ_0 and increasing ρ can effectively increase the meeting probability of the agents with different strategies, which makes the cooperators easy to be exploited by the defectors and accordingly leads to the decrease of cooperation.

4. Theoretical analysis

4.1. Relationship between extinction threshold r_L and average size $\overline{n}$ of interaction groups

In the present model, due to the purposive movement and free-floating of the agents, the interaction partners in the competing group vary with time, which will lead to the similar results in well-mixed populations where all the agents are possible to be chosen as interaction partners in the competing process. In the following, we make a mean field analysis of the replicator dynamics and give an approximation of the extinction threshold r_L in the present model.

According to the payoff function, in a randomly chosen group, a defector always gets a higher payoff than a cooperator. But in mean field analysis, the system does not evolve according to such payoffs but the averaged payoffs for cooperators or defectors which are obtained by averaging over all groups. In the present model, not all the interaction groups have the same size. In theoretical analysis, we take the average size of the interaction groups as the interaction group size, which satisfies $\overline{n} = 9\rho$.

Suppose in the well-mixed population, the frequencies of cooperators, defectors and loners are f_C, f_D and f_L respectively, which satisfy the condition $f_C + f_D + f_L = 1$. As that in ref.[17], the average payoffs for defectors, cooperators and loners are

$$P_D = \sigma f_L^{\overline{n}-1} + r\frac{f_C}{1-f_L}[1 - \frac{1-f_L^{\overline{n}}}{\overline{n}(1-f_L)}], \tag{7}$$

$$P_C = P_D - (r-1)f_L^{\overline{n}-1} + \frac{r}{\overline{n}}\frac{1-f_L^{\overline{n}}}{1-f_L} - 1, \tag{8}$$

$$P_L = \sigma. \tag{9}$$

As the multiplication rate r increases to the extinction threshold $r = r_L$, the loners become extinct and we get $f_L = 0$ and $f_C + f_D = 1$. In such a case, the payoffs for defectors and cooperators become

$$P_D = \frac{\overline{n}-1}{\overline{n}} r_L f_C, \tag{10}$$

$$P_C = \frac{\overline{n}-1}{\overline{n}} r_L f_C + \frac{1}{\overline{n}} r_L - 1. \tag{11}$$

On the condition that $P_C = P_D$, where cooperators and defectors can coexist, we obtain the extinction threshold

$$r_L = \overline{n}. \tag{12}$$

According to the above equation, the extinction threshold r_L in the present model is only related to the average size of interaction groups. The rise of population density will lead to the rise of the average group size and thereafter the increase of the extinction threshold. For example, as population density ρ increases from $\rho = 0.3$ to $\rho = 0.8$, we can estimate that r_L should increase from 2.7 to 7.2. Compared with the simulation results in fig.1, it is found that the theoretical analysis is in accordance with simulation data.

4.2. Relationship between free-floating probability ξ_0 and levels of cooperation

In simulation results, it is observed that the free-floating of the agents has great impact on the change of cooperation. In the following, by theoretical analysis, we will give a picture of what may be the possible reasons for the occurrence of such an impact.

In the OPGG, the frequencies of cooperators, defectors and loners are determined by the payoffs of the strategies. In well-mixed cases, the payoff for each agent is determined by the group size and the status of the agents in the same group. The larger the group size, the more possible for two agents to meet each other at a time. Just like that in well-mixed cases, in a mobile environment, although all the agents are on the sites of a network, they are possible to meet each other within a period of time. The faster the free-floating of the agents, the more possibly the agents meet each other within the period of time. Therefore, from the view point of the meeting probability of two agents, the effect of increasing the speed of free-floating is the same as the effect of increasing the group size. In the following, we will firstly give a functional relation between the group size and the free-floating probability, and then give a theoretical analysis of the relationship between the free-floating probability ξ_0 and levels of cooperation.

In the present model, because all the agents are arranged on the sites of a rigid network with Moore neighborhood, the number of agents in the same group should be less than or equal to 9. Considering the effects of increasing the group size and the speed of free-floating on the increase of meeting probability, we define the boundary conditions of ξ_0 and n as: for $\xi_0 \sim 0$, $n \sim 9\rho$; for $\xi_0 \sim 1$, $n \sim 9$. Therefore, the following functional relation between n and ξ_0 is adopted,

$$n(\rho, \xi_0) = 9[\rho + (1-\rho)^{\frac{1}{\xi_0}}]. \tag{13}$$

Just like that in a well-mixed case[17, 41], for predefined ρ and ξ_0, the difference of the payoff between a defector and a cooperator satisfies the equation

$$P_D - P_C = 1 + (r-1)f_L^{n(\rho,\xi_0)-1} - \frac{r}{n(\rho,\xi_0)} \frac{1 - f_L^{n(\rho,\xi_0)}}{1 - f_L}. \tag{14}$$

In the steady state, the payoffs of all the agents should be the same, $P_C = P_D = P_L = \sigma$. Therefore, the above equation becomes

$$1+(r-1)f_L^{n(\rho,\xi_0)-1}-\frac{r}{n(\rho,\xi_0)}\frac{1-f_L^{n(\rho,\xi_0)}}{1-f_L}=0. \tag{15}$$

For $0 < f_L < 1$, the solution of the above equation is the same as the solution of the following equation

$$(1-f_L)+(r-1)(1-f_L)f_L^{n(\rho,\xi_0)-1}-\frac{r}{n(\rho,\xi_0)}(1-f_L^{n(\rho,\xi_0)})=0. \tag{16}$$

Suppose

$$F(f_L,n(\rho,\xi_0))=(1-f_L)+(r-1)(1-f_L)f_L^{n(\rho,\xi_0)-1}-\frac{r}{n(\rho,\xi_0)}(1-f_L^{n(\rho,\xi_0)}). \tag{17}$$

For the three cases of $f_L = 0$, $f_L = 1$ and $f_L \to 1$, we obtain

$$F(0,n(\rho,\xi_0))=1-\frac{r}{n(\rho,\xi_0)}, \tag{18}$$

$$F(1,n(\rho,\xi_0))=0, \tag{19}$$

$$F(f_L,n(\rho,\xi_0))\approx(1-r)(n(\rho,\xi_0)-1)(1-f_L)^2. \tag{20}$$

From the equations of (18), (19) and (20) we find that, within the range of $r > 2$ and $n(\rho, \xi_0) > r$, as f_L changes from 0 to 1, the value of $F(f_L, n(\rho, \xi_0))$ changes from $F(f_L, n(\rho, \xi_0)) > 0$ to $F(f_L, n(\rho, \xi_0)) < 0$. As we check the sign of $\frac{\partial^2 F(f_L,n)}{\partial f_L^2}$, we find that it changes at most once within the range of $0 < f_L < 1$. Such results indicate that, within the range of $0 < f_L < 1$, there exists a single solution f_L^{tr} for equation (16). For $f_L > f_L^{tr}$, the payoff for cooperators is less than the payoff for defectors. For $f_L < f_L^{tr}$, the payoff for cooperators is greater than the payoff for defectors.

The above theoretical analysis indicates that, in the present model, increasing ξ_0 will lead to the increase of f_L and accordingly the decrease of f_C, which is in accordance with the simulation results found in Fig. 3.

5. Summary

In the face of the choice to win everything or get nothing in a public goods game, it is a reasonable choice for the individuals to drop out from the game and enjoy a small but fixed gain, which yields the rock-scissors-paper cycles of different strategies in the optional public goods game. The oscillatory persistence of cooperation in the optional public goods game displays the sustain-ability of cooperation, but it does not tell us on which conditions the system will evolve to slow oscillations and different levels of stable cooperation can be reached. By incorporating population density and individual mobility into the original OPGG, it is found that both the stableness and the improvement of cooperation are connected to the degree of crowdedness and the speed of free-floating of competitive agents.

For low population density and slow free-floating of competitive agents, the departure of unsatisfactory agents from competing groups make it easy for scattered cooperators to agglomerate into independent cooperator clusters which results in the expansion of cooperation. The agents who are left on the original sites are more possible to form a single defector cluster, which will finally be doomed by the random-moving loners. For low population density and fast free-floating of competitive agents, the exchange of neighbors often takes place and the defectors are easy to invade the independent clusters of cooperators. Numerous small components where cooperators and defectors coexist occur and the levels of cooperation decrease. For high population density, because of the percolation effect, nearly all the agents are merged into the same component. The defectors uniformly expand into cooperator territory and the evolutionary dynamics in the original OPGG is recovered. The relationship between the levels of cooperation and the free-floating probability is found.

The present model indicates that the existence of different levels of cooperation in the real world should be related to the environmental conditions, including population density and individual mobility. In the future, similar environmental conditions should be considered in the game models with continuous strategy variables and the generalization of these conditions is a special interest of ours.

Acknowledgments

This work is the research fruits of the Humanities and Social Sciences Fund sponsored by Ministry of Education of China (Grant No. 10YJAZH137), Natural Science Foundation of Zhejiang Province (Grant No. Y6110687), Social Science Foundation of Zhejiang Province (Grant No. 10CGGL14YB) and National Natural Science Foundation of China (Grant Nos. 10805025, 11175079, 70871019, 71171036, 71072140).

References

[1] Van Segbroeck S, Santos FC, Lenaerts T, Pacheco JM. Reacting differently to adverse ties promotes cooperation in social networks. Phys Rev Lett 2009; 102(5): 058105.

[2] Graser O, Xu C, Hui PM. Analytic approach to co-evolving dynamics in complex networks: dissatisfied adaptive snowdrift game. New J Phys 2011; 13(8): 083015.

[3] Sigmund K, De Silva H, Traulsen A, Hauert C. Social learning promotes institutions for governing the commons. Nature 2010; 466: 861-863.

[4] Perc M. Sustainable institutionalized punishment requires elimination of second-order free-riders. Scientific Reports 2012; 2:344.

[5] Wang WX, Yang R, Lai YC. Cascade of bankruptcy and emergence of pure cooperation in evolutionary games on networks. Phys Rev E 2010; 81: 035102.

[6] Yun CK, Masuda N, Kahng B. Diversity and critical behavior in prisoner's dilemma game. Phys Rev E 2011; 83: 057102.

[7] Jiang LL, Zhao M, Yang HX, Wakeling J, Wang BH, Zhou T. Reducing the heterogeneity of payoffs: an effective way to promote cooperation in prisoner's dilemma game. Phys Rev E 2009; 80: 031144.

[8] Chen YZ, Huang ZG, Wang SJ, Zhang Y, Wang YH. Diversity of rationality affects the evolution of cooperation. Phys Rev E 2009; 79: 055101.

[9] Zhong LX, Zheng DF, Zheng B, Xu C, Hui PM. Networking effects on cooperation in evolutionary snowdrift game. Europhys Lett 2006; 76(4): 724.

[10] Li PP, Ke JH, Lin Z, Hui PM. Cooperative behavior in evolutionary snowdrift games with the unconditional imitation rule on regular lattices. Phys Rev E 2012; 85: 021111.

[11] Wang WX, Ren J, Chen GR, Wang BH. Memory-based snowdrift game on networks. Phys Rev E 2006; 74: 056113.

[12] Zhong LX, Qiu T, Shi YD. Limitation of network inhomogeneity in improving cooperation in coevolutionary dynamics. Physica A 2012; 391: 2322-2329.

[13] Zheng DF, Yin HP, Chan CH, Hui PM. Cooperative behavior in a model of evolutionary snowdrift games with N-person interactions. Europhys Lett 2007; 80: 18002.

[14] Santos FC, Santos MD, Pacheco JM. Social diversity promotes the emergence of cooperation in public goods games. Nature 2008; 454: 213-217.

[15] Perc M, Szolnoki A. Self-organization of punishment in structured populations. New J Phys 2012; 14: 043013.

[16] Zhang JL, Zhang CY, Chu TG. The evolution of cooperation in spatial groups. Chaos, Solitons and Fractals 2011; 44: 131-136.

[17] Hauert C, De Monte S, Hofbauer J, Sigmund K. Volunteering as red queen mechanism for cooperation in public goods game. Science 2002; 296: 1129-1132.

[18] Rand DG, Nowak MA. The evolution of antisocial punishment in optional public goods games. Nat Commun 2011; 2:434.

[19] Xu ZJ, Wang Z, Song HP, Zhang LZ. Self-adjusting rule in spatial voluntary public goods games. Europhys Lett 2010; 90: 20001.

[20] Sasaki T, Okada I, Unemi T. Probabilistic participation in public goods games. Proc R Soc B 2007; 274: 2639-2642.

[21] Johnson NF. Financial systems: Ecology and economics. Nature 2011; 469: 302.

[22] Zhong LX, Qiu T, Ren F, Li PP, Chen BH. Time scales of epidemic spread and risk perception on adaptive networks. Europhys Lett 2011; 94: 18004.

[23] Lu LY, Zhang YC, Yeung CH, Zhou T. Leaders in social networks, the delicious case. PLoS One 2011; 6: e21202.

[24] Xu XL, Qu YQ, Guan S, Jiang YM, He DR. Interconnecting bilayer networks. Europhys Lett 2011; 93: 68002.

[25] Szabo G, Hauert C. Phase transitions and volunteering in spatial public goods games. Phys Rev Lett 2002; 89: 118101.

[26] Amritkara RE, Hu CK. Synchronized state of coupled dynamics on time-varying networks. Chaos 2006; 16: 015117.

[27] Cressman R, Vickers GT. Spatial and density effects in evolutionary game theory. J Theor Biol 1997; 184: 359-369.

[28] Rankin DJ. Resolving the tragedy of the commons: the feedback between intraspecific conflict and population density. J Evol Biol 2007; 20: 173-180.

[29] Wang Z, Szolnoki A, Perc M. Percolation threshold determines the optimal population density for public cooperation. Phys Rev E 2012; 85: 037101.

[30] Wakano JY, Nowak MA, Hauert C. Spatial dynamics of ecological public goods. Proc Natl Acad Sci USA 2009; 106: 7910-7914.

[31] Gonzalez MC, Lind PG, Herrmann HJ. System of mobile agents to model social networks. Phys Rev Lett 2006; 96: 088702.

[32] Peruani F, Sibona G. Dynamics and steady states in excitable mobile agent system. Phys Rev Lett 2008; 100: 168103.

[33] Song CM, Qu ZH, Blumm N, Barabasi AL. Limits of predictability in human mobility. Science 2010; 327: 1018-1021.

[34] Cardillo A, Meloni S, Gomez-Gardenes J, Moreno Y. Velocity-enhanced cooperation of moving agents playing public goods games. Phys Rev E 2012; 85: 067101.

[35] Helbing D, Yu WJ. The outbreak of cooperation among success-driven individuals under noisy conditions. Proc Natl Acad Sci USA 2009; 106: 3680-3685.

[36] Meloni S, Buscarino A, Fortuna L, Frasca M, Gomez-Gardenes J, Latora V, Moreno Y. Effects of mobility in a population of prisoner's dilemma players. Phys Rev E 2009; 79: 067101.

[37] Droz M, Szwabinski J, Szabo G. Motion of influential players can support cooperation in prisoner's dilemma. Eur Phys J B 2009; 71: 579-585.

[38] Perc M, Szolnoki A. Coevolutionary games-a mini review. Biosystems 2010; 99:109-125.

[39] Liu ZH. Effect of mobility in partially occupied complex networks. Phys Rev E 2010; 81: 016110.

[40] Roca CP, Helbing D. Emergence of social cohesion in a model society of greedy, mobile individuals. Proc Natl Acad Sci USA 2011; 108: 11370-11374.

[41] C. Hauert, S. De Monte, J. Hofbauer, K. Sigmund. Replicator dynamics for optional public good games. J. Theor. Biol. 2002; 218: 187-194.